

A Helch

✓

Heber

Exchange

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Abbe Jr. Hatch

{ Geo. Leo

{ Emma Sister

Drug

Store

West side

Main

Valley

Drug

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at